arXiv:1302.7126v1 [physics.soc-ph] 28 Feb 2013

Growing multiplex networks

V. Nicosia,[1] G. Bianconi,[1] V. Latora,[1] and M. Barthelemy[2]
[1]*School of Mathematical Sciences, Queen Mary University of London, Mile End Road, E1 4NS, London (UK)*
[2]*Institut de Physique Théorique, CEA, CNRS-URA 2306, F-91191, Gif-sur-Yvette, France*

We propose a modelling framework for growing multiplexes where a node can belong to different networks. We define new measures for multiplexes and we identify a number of relevant ingredients for modeling their evolution such as the coupling between the different layers and the arrival time distribution of nodes. The topology of the multiplex changes significantly in the different cases under consideration, with effects of the arrival time of nodes on the degree distribution, average shortest paths and interdependence.

PACS numbers: 89.75.Fb, 89.75.Hc and 89.75,-k

Many systems are structured as networks, and their properties are now, after a decade of efforts, well understood [1–4]. However, a complex network is rarely isolated, and some of its nodes could be part of many graphs, at the same time. Examples include multimodal transportation networks [6, 7], climatic systems [8], economic markets [9], energy-supply networks [10] and the human brain [11]. In these cases, each network has to be considered as being a part of a larger system in which a set of interdependent networks with different structure and function coexist, interact and coevolve. So far network scientists have investigated these systems by looking at one type of relationship at a time, e.g. by analyzing collaboration networks and email communications as separate graphs. However, the structural properties of each of these networks and their evolution can depend in a non-trivial way on that of other graphs to which they are interconnected. For instance, two people can become friends after having been in touch by email for a while, or two researchers start a collaboration after having attended the same conference (i.e. being part of the same co-location network for a certain amount of time). Consequently, these systems are better represented as *multiplexes*, i.e. graphs composed by M different layers in which the same set of N nodes can be connected to each other by means of links belonging to M different classes or types.

The characterization of multiplexes is still in its infancy. Some recent works have proposed suitable extensions to multi-layer graphs of classic network metrics and models [13, 17], and preliminary results show that multiplexicity has important consequences for the dynamics of processes occurring in real systems, including routing [12, 13], diffusion [14], cooperation [16] and epidemic spreading [15].

In this Letter we propose and study a generic model of multiplex growth, inspired by classical preferential-attachment based models, in which the probability for a newly arrived node to establish connections to existing nodes in each of the layers of a multiplex is a function of the degree of other nodes at all layers. We define two new metrics to characterize the structure of multiplexes, namely the inter-layer joint-degree distribution and the node interdependence, and we study the effect of different attachment rules on the structure of the resulting network. We then focus on 2-layer multiplexes, and we provide closed forms for both the degree distributions at each layer and the inter-layer degree-degree correlations corresponding to different attachment rules. We also investigate the interplay between the attachment rule and the relative time-scales of edge creation at each layer, and we show that the presence of delays in the arrival of nodes at different layers can have a significant impact on the distribution of shortest paths and interdependence.

More precisely, a multiplex is a set of N nodes which are connected to each other by means of edges belonging to M different classes or types. We represent each class of edges as a separate *layer*, and we assume that a node i of the multiplex consists of M replicas, one for each layer. We denote by $V^{[\alpha]}$ the set of nodes in layer α and by $E^{[\alpha]}$ the set of all the edges of a given type α. Consequently, the α^{th} layer of a multiplex is represented by a graph $G^{[\alpha]}(V^{[\alpha]}, E^{[\alpha]})$, associated to the adjacency matrix $A^{[\alpha]} = \{a_{ij}^{[\alpha]}\}$, where $a_{ij}^{[\alpha]} = 1$ if node i and node j are connected by an edge of type α, whereas $a_{ij}^{[\alpha]} = 0$ otherwise. An M-layer multiplex is fully specified by the vector $\mathcal{A} = [A^{[1]}, A^{[2]}, \ldots, A^{[M]}]$, whose elements are the adjacency matrices of the M layers. We denote by $k_i^{[\alpha]} = \sum_j a_{ij}^{[\alpha]}$ the degree of node i at layer α, i.e. the number of edges of type α of which i is an endpoint, and by $\boldsymbol{k}_i$ the M-dimensional vector of the degrees of the replicas of i. In general, the degrees of the replicas of i are distinct, and some replicas can also be isolated (i.e. $k_i^{[\alpha]} = 0$ for some value of α). In the following we consider all the edges at all layers to be undirected and unweighted.

As in the case of classical 'singlex' graphs, we can characterize each layer α of a multiplex by studying the degree distribution $P(k^{[\alpha]})$, and and the joint-degree distribution $P(k^{[\alpha]}, k^{[\alpha']})$. However, we are interested here in the structural properties of the multiplex as a whole, so we propose to quantify the correlations between the degrees of replicas of the same node at two different layers α

and α', by constructing the inter-layer joint-degree distributions $P(k^{[\alpha]}, k^{[\alpha']})$, or equivalently the conditional degree distributions $P(k^{[\alpha']}|k^{[\alpha]})$. In particular, we can look at the projection of the conditional distribution obtained by considering the average degree $\bar{k}^{[\alpha']}$ at layer α' of nodes having degree $k^{[\alpha]}$ at layer α:

$$\bar{k}^{[\alpha']}(k^{[\alpha]}) = \sum_{k^{[\alpha']}} k^{[\alpha']} P(k^{[\alpha']}|k^{[\alpha]}) \tag{1}$$

By plotting this quantity as a function of $k^{[\alpha]}$ we can detect the presence and the sign of degree correlations between the two layers. For a multiplex with no inter-layer correlations between layers α and α' we expect $\bar{k}^{[\alpha']}(k^{[\alpha]}) = \langle k^{[\alpha']} \rangle$ and $\bar{k}^{[\alpha]}(k^{[\alpha']}) = \langle k^{[\alpha]} \rangle$. If $\bar{k}^{[\alpha']}(k^{[\alpha]})$ increases with $k^{[\alpha]}$ we say that the degrees of the two layers have positive (assortative) correlations, while if $\bar{k}^{[\alpha']}(k^{[\alpha]})$ is a decreasing function of $k^{[\alpha]}$ we say that the degrees on layer α and α' are anticorrelated (or disassortatively correlated).



FIG. 1: (color online) **Linear attachment with synchronous arrival** Panel a-b: degree distribution $P(k)$ (left) and the projection $\bar{k}(k)$ of the inter-layer degree-degree correlations (right) for the $\{1, 0, 0, 1\}$ (top) and the $\{0, 1, 1, 0\}$ (bottom) attachment kernels (red circles are for the first layer, yellow squares for the second layer). These results correspond to 2-layer multiplexes with $N = 10000$ nodes, $m = 3$, $m_0 = 3$, and are averaged over 50 realizations. The observed degree distributions and the inter-layer degree correlations closely follow the theoretical curves (reported as solid black lines) and are relatively insensitive to the coupling matrix.

In addition to the assortativity, we can also characterize for each node i of a multiplex, its 'multiplex reachability' by computing the average distance L_i from i to any other node of the the multiplex and comparing this average distance with that measured on each layer separately. However, the presence of more than one layer in a multiplex produces an increase in the number of available paths, so that the distance between two nodes of a multiplex will be, in general, smaller than or at most equal to that measured on each layer separately. A measure to quantify the value added by the multiplexicity to the reachability of nodes is the *interdependence* [13] which for a node i is defined by

$$\lambda_i = \sum_{\substack{j \in N \\ j \neq i}} \frac{\psi_{ij}}{\sigma_{ij}} \tag{2}$$

where ψ_{ij} is the number of shortest paths between node i and node j which use edges lying on more than one layer, while σ_{ij} is the total number of shortest paths between i and j in the multiplex. The interdependence of a multiplex is computed as the average node interdependence $\lambda = 1/N \sum_i \lambda_i$ and is normalized in $[0, 1]$. If λ is close to zero, then most of the shortest paths among nodes lie on just one layer, while if λ is close to 1 then the majority of the shortest paths exploit more than one layer.

We now introduce the first model of growing multiplex networks. Most of the classical growing models for single-layer networks assume that new nodes arrive in the graph one by one and that each node carries m edge stubs; upon arrival, the new node creates m new connections with other existing nodes, according to a prescribed attachment rule. In that case, each node i has a unique *arrival time* t_i. In the case of multiplexes, instead, each layer can exhibit a different edge-formation dynamics, and in general the edges of the M replicas of a new node are not created at the same time. For instance, a face-to-face interaction relationship is usually established before two individuals become friend, while two locations are usually connected by a road before a direct railway line between them is constructed. Consequently, we assume that a newly arrived node has m stubs on each layer of the multiplex, but the replica of a node i on layer α connects its m stubs at a different time $t_i^{[\alpha]}$. We denote by $\boldsymbol{t}_i$ the vector of arrival times of the replicas of node i. In order to make the model analytically tractable, we make two simplifying assumptions. The first is that there exists a layer $\bar{\alpha}$ so that $t_i^{[\bar{\alpha}]} \leq t_i^{[\alpha]}\ \forall i, \forall \alpha \neq \bar{\alpha}$. This is equivalent to saying that a newly arrived node must first create its connections on layer $\bar{\alpha}$ before any of its replicas can create connections on any other layer $\alpha \neq \bar{\alpha}$. We call $\bar{\alpha}$ the *master layer*. The second assumption is that nodes arrive one by one on the master layer, at equal discrete time intervals $t = \{0, 1, \ldots, \}$. We label the nodes of a growing multiplex according to the ordering induced by their arrival on the master layer. Without loss of generality, in the following we assume that the the master layer is the first one, i.e. $\bar{\alpha} = 1$, and that the arrival times of the replicas of node i have the form

$$t_i^{[\alpha]} = T(t_i^{[1]}, \xi^{[\alpha]}(\tau)) \tag{3}$$

where T is a certain function of $t_i^{[1]}$ and of the random variable $\xi^{[\alpha]}(\tau)$. By appropriately choosing T and $\xi^{[\alpha]}(\tau)$ we can model different arrival behaviours, including *a)*



FIG. 2: (color online) **Linear attachment with delayed arrival.** Blue lines correspond to $\beta = 1.1$, red lines to $\beta = 2.0$. The top sub-panels of each panel correspond to the $\{0, 1, 1, 0\}$ coupling pattern, while the bottom sub-panels are for $\{1, 0, 0, 1\}$. Panel a): The degree $k(t)$ of the largest hub of the second layer as a function of time. $k(t)$ scales as $(t/t_0)^\delta$, where δ depends on the exponent β and approaches $\delta = 0.5$ when β increases (panel b). The value of β tunes the shape of the distribution of average shortest path length from each node to all the others (panel c) and the distribution of node interdependence (panel d). When β increases these distributions become more similar to the corresponding ones observed for synchronous arrival (shaded grey). All plots correspond to 2-layer multiplexes with $N = 10000$ nodes, $m = 3$, $m_0 = 3$, and the results are averaged over 50 realizations.

simultaneous arrival ($T = t_i^{[1]}$); *b) power-law delayed arrival* ($T = t_i^{[1]} + \xi(\tau)$ and $P(\xi = \tau) = (\beta - 1)\tau^{-\beta}$ for $\tau \geq 1$ and $\beta > 1$). Upon arrival, the newborn node i connects to m existing nodes in the master layer, according to a certain attachment rule. As in the preferential attachment models [5], we assume that the attachment probability depends on the degree of a node. However, in a multiplex the probability for node i to connect to node j on each layer α can depend not only on $k_j^{[\alpha]}$ but also on the degrees of j's replicas on the other layers

$$\Pi_{i\to j}^{[\alpha]} = \frac{F_j^{[\alpha]}(\boldsymbol{k}_j)}{\sum_l F_l^{[\alpha]}(\boldsymbol{k}_l)} \tag{4}$$

For the sake of clarity and without loss of generality, we focus in the following on 2-layers multiplexes with $\alpha = 1, 2$..

We begin with the simplest case of *linear attachment* which is the natural extension of the Barabasi-Albert model [5]. In this case, we consider that the probability for a newborn node i to connect to an existing node j on layer α is proportional to a linear combination of the degrees of j at all layers. Therefore, the attachment kernels can be expressed as

$$\begin{bmatrix} F^{[1]}[k,q] \\ F^{[2]}[k,q] \end{bmatrix} = C \begin{bmatrix} k \\ q \end{bmatrix} = \begin{bmatrix} c^{[1,1]} & c^{[1,2]} \\ c^{[2,1]} & c^{[2,2]} \end{bmatrix} \begin{bmatrix} k \\ q \end{bmatrix} \tag{5}$$

where from now on we denote the degree $k^{[1]}$ on layer 1 by k and the degree $k^{[2]}$ on layer 2 by q, i.e. $k^{[1]} = k$ and $k^{[2]} = q$. The coefficients $c^{[r,s]}$ tune the dependence of the attachment probability at layer r on the degrees of nodes at layer s. In the case of 2-layer multiplexes we can represent the set of coefficients $C = \{c^{[r,s]}\}$ using the compact notation $\{c^{[1,1]}, c^{[1,2]}, c^{[2,1]}, c^{[2,2]}\}$. The dynamics can be easily solved in mean-field and in some specific cases we can also solve analytically the master-equation and fully characterize the degree correlations within the two different layers. In the following we summarize the master-equation solution in some particularly interesting cases. First of all let us consider simultaneous arrival of the nodes in the two layers. If we set $C = \{1, 0, 0, 1\}$ then the attachment probability at each layer will depend only on the degree of the nodes in the same layer. In this case the degree distribution in the first layer reads

$$P(k) = \frac{2m(m+1)}{k(k+1)(k+2)}, \quad k > m \tag{6}$$

and the degree distribution in the second layer is identically equal. This distribution goes as $P(k) \sim k^{-\gamma}$ with $\gamma \simeq 3$. If we solve the master-equation for the multiplex evolution we obtain the analytical expression for the inter-layer joint degree probability $P(k, q)$

$$P(k,q) = \frac{2\Gamma(2+2m)\Gamma(1)\Gamma(q)\Gamma(k+q-2m+1)}{\Gamma(m)\Gamma(m)\Gamma(k+q+3)\Gamma(k-m+1)\Gamma(q-m+1)} \tag{7}$$

The average degree $\bar{k}(q)$ at layer 1 of nodes having degree q at layer 2 reads:

$$\bar{k}(q) = \frac{m(q+2)}{1+m} \tag{8}$$

Notice that even if the two layers grow independently, the simultaneous arrival introduces non-trivial inter-layer degree correlations. In fact, in the mean-field approach, the degree of a node on each layer increases over time as $k_i^{[\alpha]}(t) = m\left(t/t_i^{[\alpha]}\right)^{1/2}$, so that the degrees of the two replicas of a node i depend, for large t, only on their arrival time. If both replicas have the same arrival time, i.e. $t_i^{[1]} = t_i^{[2]}$ then the degree of the two replicas will be positively correlated. In Fig. 1 we report the degree distribution and the values of $\bar{k}(q)$ for two coupling patterns, which are in good agreement with the theoretical curves. It is clear from the figure that in the synchronous arrival case the shape of the coupling matrix is actually not very relevant and that the value of the degree distribution exponent and strong assortativity are robust features of these multiplexes.

If we consider a power-law delayed arrival time on the second layer, the results are significantly different. In Fig. 2 we illustrate how the exponent of the delay distribution β affects the structure of the obtained multiplex. In particular, the coupling matrix has now an effect on



FIG. 3: (color online) **Semi-linear attachment.** Degree distributions (panel a) and inter-layer degree-degree correlations (panel b) for a multiplex with semi-linear attachment and different coupling patterns, respectively $\{1,0,0,1\}$ (left) ,$\{0,1,1,0\}$ (middle) and $\{1,0,1,0\}$ (right). Red circles are for the first layer and yellow squares for the second layer. Panel c): the distribution of the average shortest path length from one node to all the other nodes heavily depends on the coupling pattern. Similarly, the interdependence of a node $\lambda(t)$ is always a sublinear function of the arrival time t but its shape depends on the coupling pattern at work (panel d). In general, older nodes have smaller interdependence. Panel e): The coupling pattern also affects the distribution of node interdependence. The smallest average interdependence is observed when the two layers are independent, i.e. case $C = \{1,0,0,1\}$ (yellow curve).

various quantities. The degree distribution are still power laws but the exponent depends now on β: for β small, some nodes become predominant thanks to the absence of other nodes and the degree distribution becomes broader. This effect being stronger for an off-diagonal matrix for which the interlayer coupling is strong. The average shortest path and the interdependence are also strongly afftected: the presence of more predominant 'old' hubs lower the average shortest path and the interlayer assortativity.

So far, we have considered the case of two scale-free growing networks and it would be interesting to capture the possibility of having a scale-free network coupled to a network with peaked degree distribution. In this respect, we introduce the *semi-linear attachment* which allows to grow multiplexes in which the two layers have different topological structures. The model is defined as follows

$$\begin{bmatrix} F^{[1]}[k,q] \\ F^{[2]}[k,q] \end{bmatrix} = C \begin{bmatrix} k \\ 1 \end{bmatrix} \tag{9}$$

where C is still a 2×2 matrix of coefficients, as in the linear model. In this case, the degree of a node on any of the two layers could depend only on its degree on layer 1 and does not ever depend on its degree on layer 2. If we set $c^{[1,1]} = c^{[2,2]} = 1$ and $c^{[1,2]} = c^{[2,1]} = 0$ we can analytically solve the master equation and the degree distributions of the two layers read:

$$P^{[1]}(k) = \frac{2m(m+1)}{k(k+1)(k+2)} \tag{10}$$

$$P^{[2]}(q) = \frac{1}{1+m}\left(\frac{m}{1+m}\right)^{q-m} \tag{11}$$

while the inter-layer joint degree distribution is equal to

$$P(k,q) = a(k)\sum_{n=0}^{k-m}\binom{k-m}{n}\left(\frac{2m}{2+2m+k-n}\right)^{q-m+1}(-1)^{k-m+n} \tag{12}$$

where $a(k) = \frac{\Gamma(k)}{\Gamma(m+1)\Gamma(k-m+1)}$. The function $\bar{k}(q)$ is given by

$$\bar{k}(q) = m\left(\frac{2(m+1)}{1+2m}\right)^{q-m+1} \tag{13}$$

Similar relations can be derived for the other coupling patterns. In Panel a) and b) of Fig. 3 we report the degree distribution and the value of $\bar{k}(q)$ for three different coupling patterns, which are in good agreeement with the theoretical curves. It is worth noticing that in the semi-linear model the coupling pattern has a dramatic impact on other structural properties of the multiplex such as the distribution of the average shortest path length from each node and the distribution of node interdependence. In particular, the interdependence is larger for older nodes, and grows sublinearly with time. This implies in particular, that navigation for old nodes is easier within a single layer while younger nodes will have to resort to the different layers to reach a target. In addition, the growth being sublinear the performance of the system increases very slowly.

In summary, we have presented a framework for modelling the growth of multiplexes which enabled us to highlight the relevant ingredients for these systems and which, we believe, will provide a solid basis for more elaborated models.

VN and VL acknowledge support from the EU-LASAGNE Project, Contract No.318132 (STREP) funded by the European Commission. MB thanks Queen Mary University for its warm welcome at the start of this project, and is supported by the FET-Proactive project PLEXMATH (FP7-ICT-2011-8; grant number 317614) funded by the European Commission.

[1] R. Albert and A.-L. Barabasi, *Reviews of Modern Physics* **74,**, 47 (2002).

[2] M. E. J. Newman, *SIAM Review* **45,**, 167-256 (2003).

[3] S. Boccaletti, V. Latora, Y. Moreno, M. Chavez and D.-U. Hwang, *Physics Reports* **424**, 175 - 308 (2006).

[4] A. Barrat, M. Barthelemy and A. Vespignani, , (2008).

[5] R. Albert, H. Jeong and A.-L. Barabasi, *Nature* **401**, 130–131 (1999).

[6] M. Kurant and P. Thiran, *Phys. Rev. Lett.* **96**, 138701 (2006).

[7] S.-R. Zou, T. Zhou, A.-F. Liu, X.-L. Xu and D.-R. He, *Physics Letters A* **374**, 4406 - 4410 (2010).

[8] J. Donges, H. Schultz, N. Marwan, Y. Zou and J. Kurths, *The European Physical Journal B* **84**, 635-651 (2011).

[9] J. Yang, W. Wang and G. Chen, *Physica A: Statistical Mechanics and its Applications* **388**, 2435 - 2449 (2009).

[10] S. V. Buldyrev, R. Parshani, G. Paul, H. E. Stanley and S. Havlin, *Nature* **464**, 1025–1028 (2010).

[11] E. Bullmore and O. Sporns, *Nat Rev Neurosci* **10**, 186–198 (2009).

[12] Y. Zhuo, Y. Peng, C. Liu, Y. Liu and K. Long, *Physica A: Statistical Mechanics and its Applications* **390**, 2401 - 2407 (2011).

[13] R. G. Morris and M. Barthelemy, *Phys. Rev. Lett.* **109**, 128703 (2012).

[14] S. Gómez, A. Díaz-Guilera, J. Gómez-Gardeñes, C. J. Pérez-Vicente, Y. Moreno and A. Arenas, *Phys. Rev. Lett.* **110**, 028701 (2013).

[15] A. Saumell-Mendiola, M. Á. Serrano and M. Boguñá, *Phys. Rev. E* **86**, 026106 (2012).

[16] J. Gomez-Gardeñes, I. Reinares, A. Arenas and L. M. Floria, *Sci. Rep.* **2**, – (2012).

[17] K.-M. Lee, J. Y. Kim, W. kuk Cho, K.-I. Goh and I.-M. Kim, *New Journal of Physics* **14**, 033027 (2012).

[18] R. Parshani, C. Rozenblat, D. Ietri, C. Ducruet and S. Havlin, *EPL (Europhysics Letters)* **92**, 68002 (2010).